Detection of Soybean derived materials by Real-Time PCR — Soybean detection method using *lectin 1 (Le1)*

Contents

1	Purpose and Scope	. 1
2	Principle	. 1
	Reagents and Materials	. 1
3.2	PCR reagents	. 2
4	Apparatus	. 2
5	Apparatus Procedure PCR setup	. 2
5 5.1	Procedure	. 2 . 2

1 Purpose and Scope

This document describes a taxon-specific quantitative Real-Time PCR method for the determination of the presence of soybean taxon-specific DNA. The method amplifies a 102 base pairs sequence of the soybean *lectin 1 (Le1)* gene (GenBank accession number K00821.1 M30884). This method is intended for use by an operator trained in performing Real-Time PCR analyses.

The method, when validated, may be applicable to DNA isolated from seed and grain containing mixtures, plant material, and food and feed. Its primary use is as a comparator in quantitative determination of transgenic soybean in seed and grain but it also can be used to show that DNA is of suitable quality or quantity for a PCR reaction. DNA can be extracted using a suitable DNA extraction method if the method is validated for the matrix before use. The DNA to be analyzed should be tested for quality and quantity prior to the use in the Real-Time PCR assay. Before use, the DNA extraction and PCR method should be subjected to an in-house validation procedure (ISO 17025).

2 Principle

Detection of the soybean taxon-specific DNA sequence of the *Le1* gene is performed in a in a real-time PCR using a forward and reverse primer and a *Le1* gene specific probe labeled with JOE fluorescent reporter and an BHQ1 fluorescence quencher. The *Le1* gene is present in a single copy per haploid genome (1).

3 Reagents and Materials

3.1 General

Only chemicals and water of analytical grade, appropriate for molecular biology, should be used. Solutions should be prepared by dissolving the corresponding

reagents in water and be autoclaved unless otherwise indicated. For all operations for which gloves are used it should be ensured that these are powder-free. The use of aerosol protected pipette tips (for protection against cross contamination) is recommended.

3.2 PCR reagents

Ready-to-use PCR reagent mixtures or mixtures of individual components are typically used. It is important to avoid contamination of all buffers with DNA or DNAse enzymes. Oligonucleotides used for this method are listed in Table 1.

Table 1 — Oligonucleotides

Name	Sequence		
Forward Primer	5'- CTTTCTCgCACCAATTgACA -3'		
Reverse Primer	5'- TCAAACTCAACAgCgACgAC -3'		
Probe ^a 5'-JOE- CCACAAACACATgCAggTTATCTTgg -BHQ1-3'			
^a Equivalent reporter dyes and/or quencher dyes may be used for the probe if it can be shown to yiel			

similar or better results.

4 Apparatus

The usual laboratory equipment is required, and a Real-Time PCR device suitable for the excitation of fluorescent molecules and the detection of fluorescence signals generated during PCR.

5 Procedure

The DNA used for the reaction should be of sufficient quality to avoid inhibition of the PCR reaction.

5.1 PCR setup

The method described applies for a total reaction volume of 25 μ l per PCR. (Table 2.). Method has been previously validated (BASF), changes to method would need to be validated.

Thaw the reagents, if necessary, and handle as recommended. Prepare a PCR master mix which contains all the components except for the sample DNA. Aliquot master mix and add control or sample DNA just prior to running the PCR reaction.

Final Concentration	Volume per Reaction (µl)
N/A	Up to 25uL
1X	12.5
200 nM	0.5
200 nM	0.5
200 nM	0.5
	100-300 ng
	N/A 1X 200 nM 200 nM

Table 2 — PCR Reaction Setup ^a

^a Different stock concentration for primers and probe may be used, as long as final concentration and total reaction component ratios are maintained.

^b For example, 2x TaqMan Universal PCR Master Mix

5.2 Thermocycling parameters

Run the PCR with general cycling conditions listed below in the Real-Time PCR detection platform. Check manufacturer product bulletin for specific PCR platform compatibility and cycling parameters.

Table	3 — 0	Cycling	Program ^a	1
-------	-------	---------	----------------------	---

			Measurement	of Cycles
1 UNG Incubat	50°C	2 minutes	No	1
2 Denatu	re 95°C	10 minutes	No	1
3 Anneal a Exten		15 seconds 60 seconds	No Yes	- 40

^a Changes to the conditions included in this table are not recommended and must be validated.

6 Citations

 (1) : Identification of single target taxon-specific reference assays for the most commonly genetically transformed crops using digital droplet PCR – Jacchia S. et al

Method provided by BASF, Research Triangle Park, NC, USA